



DyNAzyme™ II

PCR Master Mix

Product codes:

F-508S, 200 reactions

F-508L, 1000 reactions

Stable for six months from the packaging date. Recommended storage temperature -20°C . After thawing the mix can be refrozen or optionally stored at $+4^{\circ}\text{C}$ for three months.

1. Introduction

Finnzymes' DyNAzyme™ II DNA Polymerase is a recombinant thermostable DNA polymerase purified from an *E. coli* strain expressing the cloned DyNAzyme DNA Polymerase gene from *Thermus brockianus*. DyNAzyme II DNA Polymerase possesses the following activities: $5' \rightarrow 3'$ DNA polymerase activity and $5' \rightarrow 3'$ exonuclease activity.

DyNAzyme II PCR Master Mix is a 2x master mix containing DyNAzyme II DNA Polymerase, nucleotides, and optimized reaction buffer including MgCl_2 . Only template and primers need to be added by the user. DyNAzyme II PCR Master Mix offers several advantages when compared to *Taq* DNA polymerase, including improved thermal stability and tolerance against DMSO. In addition, high purity level and batch-to-batch consistency ensure the accurate performance of DyNAzyme II PCR Master Mix.

2. Package information

F-508S	200 reactions in 50 μl volume 2x DyNAzyme™ II PCR Master Mix (4 x 1.25 ml) Contains: 0.04 U/ μl DyNAzyme™ II DNA Polymerase, 20 mM Tris-HCl (pH 8.8 at 25°C), 3 mM MgCl_2 , 100 mM KCl, stabilizers and 400 μM of each dNTP
F-508L	1000 reactions in 50 μl volume 2x DyNAzyme™ II PCR Master Mix (20 x 1.25 ml) Contains: 0.04 U/ μl DyNAzyme™ II DNA Polymerase, 20 mM Tris-HCl (pH 8.8 at 25°C), 3 mM MgCl_2 , 100 mM KCl, stabilizers and 400 μM of each dNTP

Material safety datasheet (MSDS) is available at www.finnzymes.com.

3. Guidelines for using DyNAzyme™ II PCR Master Mix

Carefully mix and centrifuge all tubes before opening to ensure homogeneity and improve recovery. PCR reactions should be set up on ice.

Table 1. Pipetting instructions (add items in this order).

Component	50 μl reaction	20 μl reaction	Final concentr.
H_2O	add to 50 μl	add to 20 μl	
2x DyNAzyme™ II PCR Master Mix	25 μl	10 μl	1x
primer A*	x μl	x μl	0.5 μM
primer B*	x μl	x μl	0.5 μM
template DNA	x μl	x μl	

* The recommendation for final concentration is 0.5 μM but if can be optimized in a range of 0.2–1.0 μM , if needed.

Table 2. General cycling instructions.

Cycle step	2-step protocol		3-step protocol		Cycles
	Temp.	Time	Temp.	Time	
Initial denaturation	94°C	1–2 min	94°C	1–2 min	1
Denaturation	94°C	15 s–1 min	94°C	15 s–1 min	25–35
Annealing (see 5.2)	–	–	$T_m - 5^{\circ}\text{C}$	10–30 s	
Extension	72°C	40 s/kb	72°C	40 s/kb	
Final extension	72°C 4°C	5–10 min hold	72°C 4°C	5–10 min hold	1

4. Notes about reaction components

4.1 Enzyme

In DyNAzyme II PCR Master Mix the enzyme concentration is optimized. When pipetted according to the instructions, the final concentration is 1 U of enzyme in a 50 μl reaction.

4.2 Mg^{2+} and dNTP

DyNAzyme II PCR Master Mix provides 1.5 mM MgCl_2 and 200 μM dNTP in final reaction concentration.

4.3 Template

General guidelines for low complexity DNA (e.g. plasmid, lambda or BAC DNA) are: 1 μg –10 ng per 50 μl reaction volume. For high complexity genomic DNA, the amount of DNA template should be 50–500 ng per 50 μl reaction volume. If cDNA synthesis reaction mixture is used directly as a source for the template, the volume used should not exceed 10 % of the final PCR reaction volume.

4.4 PCR additives

PCR additives such as DMSO, formamide, glycerol and betaine are compatible with DyNAzyme I DNA Polymerase. We recommend using PCR additives in the following concentrations: DMSO 2–10 %, formamide 2–10 %, glycerol 5–10 %, or combinations of these. Recommended starting point is 5 % DMSO.

Note: If high DMSO concentration is used, the annealing temperature must be decreased, as DMSO alters the melting point of the primers. It has been reported that 10 % DMSO decreases the annealing temperature by 5.5–6.0 $^{\circ}\text{C}$.¹

5. Notes about cycling conditions

5.1 Denaturation

After an initial 1–2 min denaturation at 94°C, keep the denaturation as short as possible (usually 30 seconds or less at 94°C). **Note:** The denaturation time and temperature also depend on the ramp rate and temperature control mode of the cycler.

5.2 Primer annealing

The T_m 's should be calculated with the nearest-neighbor method² as results from primer T_m calculations can vary significantly depending on the method used. Instructions for T_m calculation and a link to a calculator using the nearest-neighbor method can be found on Finnzymes' website (www.finnzymes.com). We suggest the primers to be annealed for one minute or less at the highest temperature that will permit annealing of the primers to the template. A guideline for DyNAzyme II DNA Polymerase is to use an annealing temperature 5°C below the lower T_m of the primers. Two-step cycling without an annealing step is recommended for high T_m primer pairs.

5.3 Extension

The extension should be performed at 72°C (40 seconds per kilobase of amplified product).

6. Troubleshooting

No product at all or low yield
<ul style="list-style-type: none">Repeat and make sure that there are no pipetting errors.Titrate template amount.Template DNA may be damaged. Use carefully purified template.Increase extension time.Increase the number of cycles.Decrease annealing temperature.Titrate DMSO (2–8 %) in the reaction (see section 4.4).Optimize the denaturation time.Check the condition of the primers.Check the purity and concentration of the primers.
Non-specific products - High molecular weight smears
<ul style="list-style-type: none">Shorten extension time (see section 5.3).Reduce the total number of cycles.Increase annealing temperature or try 2-step protocol (see section 5.2).Vary denaturation temperature (see section 5.1).Decrease primer concentration.
Non-specific products - Low molecular weight discrete bands
<ul style="list-style-type: none">Increase annealing temperature (see section 5.2).Shorten extension time (see section 5.3).Titrate template amount.Decrease primer concentration.Design new primers.

7. Component specifications of DyNAzyme™ II PCR Master Mix

2x DyNAzyme II PCR Master Mix contains 0.04 U/ μ l DyNAzyme II DNA Polymerase, 20 mM Tris-HCl (pH 8.8 at 25°C), 3 mM MgCl₂, 100 mM KCl, stabilizers and 400 μ M of each dNTP.

DyNAzyme II DNA Polymerase has a half life of 2.5 h at 96°C. DyNAzyme II DNA Polymerase is free of contaminating endo- and exonucleases.

DNA amplification assay: Performance in PCR is tested by the amplification of 500 bp λ DNA and 6 kb M13 DNA.

8. References

- Chester N. & Marshak D.R. (1993) *Analytical Biochemistry* 209: 284–290.
- Breslauer K.J. et al., (1986) *PNAS* 83: 3746–3750.

Shipping and storage

DyNAzyme II PCR Master Mix is shipped on gel ice. Upon arrival, store the components at -20°C. DyNAzyme II PCR Master Mix is stable for six months from the packaging date when stored and handled properly. After thawing the DyNAzyme II PCR Master Mix can be refrozen or optionally stored at +4°C for three months.

Warranty

Finnzymes Oy warrants that its products will meet the specifications stated on the technical data section of the data sheets, and Finnzymes Oy agrees to replace the products free of charge if the products do not conform to the specifications. Notice for replacement must be given within 60 days of receipt. In consideration of the above commitments by Finnzymes Oy, the buyer agrees to and accepts the following conditions:

- That this warranty is in lieu of all other warranties, express or implied;
- That **ALL WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE ARE HEREBY EXCLUDED AND WAIVED**;
- That the buyer's sole remedy shall be to obtain replacement of the product free of charge from Finnzymes Oy; and
- That this remedy is in lieu of all other remedies or claims for damages, consequential or otherwise, which the buyer may have against Finnzymes Oy.

Exclusive terms of sale

Finnzymes Oy does not agree to and is not bound by any other terms or conditions, unless those terms and conditions have been expressly agreed to in writing by a duly authorised officer of Finnzymes Oy. Prices are subject to change without notice.

Recommended guidelines for safe use of the products

Finnzymes Oy recommends that the buyer and other persons using the products follow the Guidelines for Research involving Recombinant DNA Molecules (NIH guidelines) Federal Register, July 5, 1994 (59 FR 34496) and any amendments thereto. Finnzymes Oy disclaims any and all responsibility for any injury or damage which may be caused by the failure of the buyer or any other person to follow said guidelines.

Research use only

Since these products are intended for research purposes by qualified persons, the Environmental Protection Agency does not require us to supply Premanufacturing Notice.

Notice to user

The information presented here is accurate and reliable to the best of our knowledge and belief, but is not guaranteed to be so. Nothing herein is to be construed as recommending any practice or any product in violation of any patent or in violation of any law or regulation. It is the user's responsibility to determine for himself or herself the suitability of any material and/or procedure for a specific purpose and to adopt such safety precautions as may be necessary.

DyNAzyme™ is a trademark of Finnzymes Oy.

The quality system of Finnzymes Oy is certified according to standard SFS-EN ISO9001:2000.

Version 1.4, March 2009