



FINNZYMES

DyNAzyme™ EXT

PCR Kit

Product codes:

F-552S, 200 U

F-552L, 1 000 U

Stable for one year from the packaging date. Store at -20°C.

1. Introduction

Finnzymes' DyNAzyme™ EXT DNA Polymerase is a versatile and easy to use enzyme, with powerful advantages for all PCR applications. It is especially suitable for difficult templates, long PCR, and cloning. DyNAzyme EXT DNA Polymerase is an optimized mixture of DyNAzyme II DNA Polymerase and a proofreading enzyme. The DNA polymerase activity of DyNAzyme II DNA Polymerase provides the highly efficient polymerization needed for successful long and difficult PCR and also for high yields in standard PCR. The proofreading activity removes misincorporated nucleotides, which would otherwise either be extended and thus produce mutations, or block further chain extension.

DyNAzyme EXT DNA Polymerase possesses the following activities: 5'→3' DNA polymerase activity, 5'→3' exonuclease activity, and a weak 3'→5' exonuclease activity. It is capable of adding a non-templated adenine residue at the 3' end of a DNA fragment.

DyNAzyme EXT PCR Kit includes λ DNA control template and primers for 500 bp and 20 kb amplicons. The control template and primer amounts are sufficient for performing 20 amplifications with the 500 bp primers or 20 amplifications with the 20 kb primers.

2. Kit components

Component	Concentration	F-552S	F-552L
DyNAzyme™ EXT DNA Polymerase	1 U/μl	200 U	2 x 500 U
10x Optimized DyNAzyme™ EXT Buffer*		1 x 1.5 ml	4 x 1.5 ml
10x Mg ²⁺ -free DyNAzyme™ EXT Buffer		1 x 1.5 ml	4 x 1.5 ml
dNTP mix	10 mM each	2 x 100 μl	2 x 500 μl
MgCl ₂ solution	50 mM	1 x 1.5 ml	2 x 1.5 ml
Lambda control template	0.5 ng/μl	40 μl	40 μl
500 bp control primer mix	25 μM each	20 μl	20 μl
20 kb control primer mix	25 μM each	20 μl	20 μl
Ready to use DNA standard	100 ng/μl	400 μl	400 μl
Gel loading dye		1 ml	1 ml
DMSO	100 %	0.5 ml	0.5 ml

* 10x Optimized DyNAzyme EXT Buffer provides 1.5 mM MgCl₂ in final reaction concentration.

3. Guidelines for using DyNAzyme EXT DNA Polymerase

3.1 Standard and long PCR

Optimized DyNAzyme EXT Buffer (1x buffer contains 1.5 mM Mg²⁺) is recommended for standard and long PCR reactions up to 10 kb. Some applications may require lower Mg²⁺ concentrations. In these cases, use the Mg²⁺-free DyNAzyme EXT Buffer with 50 mM MgCl₂ solution provided. For optimization of especially long extension reactions, the magnesium concentration in either buffer can be supplemented using the 50 mM MgCl₂ solution provided. See section 6.2 for more information.

3.2 Difficult templates

Often the basic reaction conditions are sufficient for successful amplification of difficult templates, such as GC-rich and looped sequences. The most difficult PCR reactions can be improved by using DyNAzyme EXT DNA Polymerase together with a PCR additive, e.g. DMSO, formamide, glycerol, or betaine, which relax DNA, thus making template denaturation easier. A recommended starting point is 5 % DMSO. See section 6.4 for more information on PCR additives.

3.3 Cloning

The improved fidelity and robustness of DyNAzyme EXT DNA Polymerase make it a successful enzyme for PCR cloning. DyNAzyme EXT DNA Polymerase is capable of adding a non-templated adenine residue at the 3' end of a DNA fragment. PCR products produced with DyNAzyme EXT DNA Polymerase can be used in either TA cloning or blunt cloning.

4. Basic reaction conditions for PCR

Mix and centrifuge all tubes before opening to improve recovery. Store the tubes on ice. Always pipette DyNAzyme EXT DNA Polymerase carefully and gently. The high glycerol content (50 %) in the storage buffer may otherwise lead to pipetting errors.

Depending on the length and complexity of your starting template, optimal reaction conditions with DyNAzyme EXT DNA Polymerase vary. See the suggested pipetting instructions in table 1.

Long PCR is very sensitive even to small variations in the reaction conditions. Therefore, the optimal conditions need to be determined experimentally.

Useful tips for long PCR:

- Use UTW™ (ultra-thin wall) or standard thin-walled tubes.
- Perform manual hot start or place reaction mixtures to a pre-heated cyclor block directly from ice.

Table 1. Pipetting instructions for different size and complexity amplifications.

Fragment size / Low complexity DNA (e.g. λ DNA)	<10 kb	10-20 kb	>20 kb			
Fragment size / High complexity DNA (e.g. human genomic DNA)	<7 kb	7-12 kb	> 12 kb			
Component	Volume	Final conc.	Volume	Final conc.	Volume	Final conc.
H ₂ O	Add to 50 μl		Add to 50 μl		Add to 50 μl	
10x Optimized DyNAzyme EXT Buffer	5 μl	1x (1.5 mM MgCl ₂)	-	-	-	-
10x Mg ²⁺ -free DyNAzyme EXT Buffer	-	-	5 μl	1x	5 μl	1x
50 mM MgCl ₂	-	1.5 mM	1.7 μl	1.7 mM	2.3 μl	2.3 mM
10 mM dNTPs	1 μl	200 μM each	1.8 μl	360 μM each	2.5 μl	500 μM each
Primer A	X μl	0.5 μM*	X μl	0.5 μM*	X μl	0.5 μM*
Primer B	X μl	0.5 μM*	X μl	0.5 μM*	X μl	0.5 μM*
Template DNA (see 6.3)	X μl		X μl		X μl	
(DMSO, optional) (see 6.4)	(2.5 μl)	(5 %)	(2.5 μl)	(5 %)	(2.5 μl)	(5 %)
DyNAzyme EXT DNA Polymerase (see 6.1)	0.5-3 μl	0.5-3 U	0.5-3 μl	0.5-3 U	0.5-3 μl	0.5-3 U

* The recommendation for final concentration is 0.5 μM but it can be optimized in a range of 0.2-1.0 μM, if needed.

5. Cycling conditions

Table 2. Cycling instructions for fragments <10 kb.

Cycle step	2-step protocol		3-step protocol		Cycles
	Temp.	Time	Temp.	Time	
Initial denaturation	94°C	1-2 min	94°C	1-2 min	1
Denaturation (see 7.1)	94°C	15 s-1 min	94°C	15 s-1 min	25-35
Annealing (see 7.2)	–	–	T _m - 5°C	15 s-1 min	
Extension (see 7.3)	72°C	40 s/1 kb	72°C	40 s/1 kb	
Final extension	72°C 4°C	5-10 min hold	72°C 4°C	5-10 min hold	1

Table 3. Cycling instructions for fragments ≥10 kb.

Cycle step	2-step protocol		3-step protocol		Cycles
	Temp.	Time	Temp.	Time	
Initial denaturation	94°C	1-2 min	94°C	1-2 min	1
Denaturation (see 7.1)	94°C	15 s-1 min	94°C	15 s-1 min	10
Annealing (see 7.2)	–	–	T _m - 5°C	15 s-1 min	
Extension (see 7.3)	70°C	40 s/1 kb	70°C	40 s/1 kb	
Denaturation (see 7.1)	94°C	15 s-1 min	94°C	15 s-1 min	15-20
Annealing (see 7.2)	–	–	T _m - 5°C	15 s-1 min	
Extension (see 7.3)	70°C	40 s/1 kb + 20 s/cycle	70°C	40 s/1 kb + 20 s/cycle	
Final extension	70°C 4°C	5-10 min hold	70°C 4°C	5-10 min hold	1

6. Notes about reaction components

6.1 Enzyme

The amount of enzyme depends on the amount of template and the length of the PCR product.

Usually 1 unit of DyNAzyme EXT DNA Polymerase per 50 µl reaction volume gives good results, but for difficult templates and long PCR the optimal amount can be 0.5 - 3 U per 50 µl reaction.

6.2 Mg²⁺ concentration and dNTP concentration

Optimization of Mg²⁺ is critical since DyNAzyme EXT DNA Polymerase is a magnesium dependent enzyme. In addition to DyNAzyme EXT DNA Polymerase, the template DNA, primers and dNTPs bind Mg²⁺. Therefore, the optimal Mg²⁺ concentration will depend on the dNTP concentration, the specific template DNA and the sample buffer composition.

Excessive Mg²⁺ stabilizes the DNA double strand and prevents complete denaturation of DNA, which reduces yield. Excess Mg²⁺ can also stabilize spurious annealing of primers to incorrect template sites, decreasing specificity. On the other hand, inadequate Mg²⁺ reduces the amount of product. In general, the acceptable Mg²⁺ concentration range narrows as the length of the PCR product increases. Usually the optimal Mg²⁺ concentration is 0.5 - 1 mM over the total dNTP concentration for standard PCR and 0.1 - 0.5 mM over the total dNTP concentration for long PCR.

High quality dNTPs (e.g. F-560) should be used for optimal performance with DyNAzyme EXT DNA Polymerase. The polymerase cannot read dUTP-derivatives or dITP in the template strand so the use of these analogues is not recommended.

6.3 Template

Template preparation becomes particularly important when performing long PCR. The amount of template required depends on the length of the PCR product. For longer extensions, more template is needed.

General guidelines are: 1 pg - 10 ng / 50 µl reaction with low complexity genomes (e.g. plasmid, BAC, lambda DNA); 50-500 ng / 50 µl reaction with high complexity genomes (e.g. human genomic DNA). If cDNA synthesis reaction mixture is used as a source of template, the volume of the template should not exceed 10 % of the final PCR reaction volume.

6.4 PCR additives

DyNAzyme EXT DNA Polymerase tolerates high DMSO concentrations (up to 10 %) needed for opening of difficult templates. Also other additives which help DNA denaturation can be used with DyNAzyme EXT DNA Polymerase (formamide, glycerol, betaine and combinations of these).

We recommend using PCR additives in the following concentrations: DMSO 2-10 %, formamide 2-10 %, glycerol 5-10 %, or combinations of these. Recommended starting point is 5 % DMSO.

Note: In high DMSO concentrations the annealing temperature must be lowered, because DMSO decreases the melting point of the primers. It has been reported that 10 % DMSO decreases the melting temperature by 5.5-6.0°C.¹

7. Notes about cycling conditions

7.1 Denaturation

After an initial 1–2 min denaturation at 94°C, keep the denaturation as short as possible (usually 30 seconds or less at 94°C). This is particularly important for long PCR. **Note:** The denaturation time and temperature depend also on the ramp rate and temperature control mode of the cycler.

7.2 Primer annealing

We suggest the primers to be annealed for one minute or less at the highest temperature that will permit annealing of the primers to the template. A guideline for determination of the annealing temperature is to use a temperature 5°C lower than the lower T_m calculated by the nearest-neighbor method.² For long PCR, the primers should be designed to allow a high annealing temperature for maximal reaction specificity (preferably ≥ 65°C). If the T_m's of the primers are high enough, annealing and extension can be done in a single 70–72°C step (two-step PCR).

Instructions for T_m calculation and a link to a calculator using the nearest-neighbor method can be found on Finnzymes' website (www.finnzymes.com).

7.3 Extension

The extension for standard PCR should be performed at 72°C, and for long PCR at 68-70°C. For amplification of shorter DNA fragments (< 10 kb), a constant extension time can be used (40 seconds per one kilobase). For amplification of long DNA fragments (≥ 10 kb), use a constant extension time (40 seconds per one kilobase) for the first 10 cycles. Then, during the next 15-20 cycles, add 20 seconds to the elongation time on each cycle.

8. Amplifying control templates

8.1 Amplification of the 500 bp control

Table 4. Pipetting instructions (add items in this order).

Component	Volume / 50 µl reaction	Final conc.
H ₂ O	40 µl	
10x Optimized DyNAzyme™ EXT Buffer	5 µl	1x
10 mM dNTPs	1 µl	200 µM each
500 bp control primer mix	1 µl	0.5 µM
Lambda control template	2 µl	1 ng/50 µl
DyNAzyme™ EXT DNA Polymerase	1 µl	0.02 U/µl

Table 5. Cycling protocol for 500 bp control reactions.

Cycle step	Temp.	Time	Number of cycles
Initial denaturation	94°C	1 min	1
Denaturation	94°C	20 s	25
Annealing	60°C	20 s	
Extension	72°C	20 s	
Final extension	72°C 4°C	10 min hold	1

500 bp control amplicon in DyNAzyme EXT PCR Kit can be used to check for PCR inhibitors in DNA preparations by substituting, for example, 5 µl of the water in the standard 500 bp control protocol with 5 µl of the DNA preparation with possible inhibitors. Compare the amount of the 500 bp product made in the standard reaction with the reaction where the DNA preparation with possible inhibitors was added. Greatly reduced yields can indicate the presence of PCR inhibitors in the tested DNA preparation.

The 500 bp control reaction is quite robust and can tolerate annealing temperatures over a very broad range. When running this kit's 500 bp control amplicon with your own template/primer reaction, adjust the cycling protocol to best fit the conditions for your primers.

8.2 Amplification of the 20 kb control

Table 6. Pipetting instructions (add items in this order).


Component	Volume / 50 µl reaction	Final conc.
H ₂ O	37.5 µl	
10x Mg ²⁺ -free DyNAzyme™ EXT Buffer	5 µl	1x
50 mM MgCl ₂	1.7 µl	1.7 mM
10 mM dNTPs	1.8 µl	360 µM each
20 kb control primer mix	1 µl	0.5 µM
Lambda control template	2 µl	1 ng/50 µl
DyNAzyme™ EXT DNA Polymerase	1 µl	0.02 U/µl

Note: It is important to place the reaction tubes to a pre-heated PCR block directly from ice to prevent non-specific amplification.

Table 7. Cycling protocol for 20 kb control reactions.

Cycle step	Temp.	Time	Number of cycles
Initial denaturation	94°C	1 min	1
Denaturation	94°C	30 s	10
Annealing	65°C	30 s	
Extension	68°C	14 min	
Denaturation	94°C	30 s	15
Annealing	65°C	30 s	
Extension	68°C	14 min+ 20 s / cycle	
	4°C	hold	1

8.3 Analysis of the control reactions



After cycling is complete, add 15 µl of gel loading dye (F-350), mix thoroughly and load 20 µl on an agarose gel (1% agarose in TAE buffer). Compare the results to the image on the left.

Lane 1. DNA size standard
Lane 2. 500 bp control amplicon
Lane 3. 20 kb control amplicon
Lane 4. DNA size standard

9. Troubleshooting

No product at all or low yield
<ul style="list-style-type: none"> Repeat and make sure that there are no pipetting errors. Use fresh high quality dNTPs. Do not use dNTP mix that contains dUTP or dITP. Titrate template amount. Template DNA may be damaged. Use carefully purified template. Increase extension time. Increase the number of cycles. Decrease annealing temperature. Use more enzyme. Optimize magnesium concentration. Try adding 2-10 % DMSO in the reaction. Optimize denaturation temperature. Optimize denaturation time. Check the purity and concentration of the primers. Check primer design.
Non-specific products
<ul style="list-style-type: none"> Increase annealing temperature. Use manual hot start or place reaction mixtures to a pre-heated cycler block directly from ice. Decrease primer concentration. Design new primers. Optimize magnesium concentration.
Non-specific high molecular weight smear
<ul style="list-style-type: none"> Decrease magnesium concentration, e.g. using 0.2 mM steps. Use less enzyme. Increase dNTP concentration. Shorten extension time. Reduce the number of cycles.

10. Component specifications

10.1 DyNAzyme EXT DNA Polymerase (F-505)

DyNAzyme EXT DNA Polymerase is an optimized mixture of DyNAzyme II DNA Polymerase and a proofreading enzyme. DyNAzyme II DNA Polymerase is purified from an *E.coli* strain expressing the cloned DyNAzyme DNA Polymerase gene from *Thermus brockianus*, Finnzymes' proprietary bacterial strain. DyNAzyme EXT DNA Polymerase is purified free of contaminating endo- and exonucleases. It has a half life of 3.5 h at 96°C.

Storage buffer: 20 mM Tris-HCl (pH 7.4 at 25°C), 0.1 mM EDTA, 1 mM DTT, 100 mM KCl, stabilizers, 200 µg/ml BSA and 50 % glycerol.

Unit definition: One unit is defined as the amount of enzyme that will incorporate 10 nmoles of dNTPs into acid-insoluble form at 74°C in 30 minutes under the stated assay conditions.

Unit assay conditions: Incubation buffer: 25 mM TAPS-HCl, pH 9.3 (at 25°C), 50 mM KCl, 2 mM MgCl₂, 1 mM β-mercaptoethanol, 100 µM dCTP, 200 µM each of dATP, dGTP, dTTP. Incubation procedure: 20 µg activated calf thymus DNA and 0.5 µCi [α-³²P] dCTP are incubated with 0.1 units of DNA polymerase in 50 µl incubation buffer at 74°C for 10 minutes. The amount of incorporated dNTPs is determined by trichloroacetic acid precipitation.

Exonuclease activity: Incubation of 10 U for 4 hours at 72°C in 50 µl assay buffer with 1 µg sonicated ³H labeled ssDNA (2x10⁵ cpm/µg) released < 1 % of radioactivity.

Endonuclease assay: No endonuclease activity was observed after incubation of 10 U of DNA polymerase with 1 µg of λ DNA in assay buffer at 72°C for 4 hours.

DNA amplification assay: Performance in PCR is tested by the amplification of 20 kb and 30 kb lambda DNA.

10.2 10x Optimized DyNAzyme EXT Buffer (F-514)

In final 1x reaction concentration the Optimized DyNAzyme EXT Buffer contains 50 mM Tris-HCl (pH 9.0 at 25°C), 1.5 mM MgCl₂, 15 mM (NH₄)₂SO₄ and 0.1 % Triton® X-100.

10.3 10x Mg²⁺-free DyNAzyme EXT Buffer (F-512)

In final 1x reaction concentration the Mg²⁺-free DyNAzyme EXT Buffer contains 50 mM Tris-HCl (pH 9.0 at 25°C), 15 mM (NH₄)₂SO₄ and 0.1 % Triton® X-100.

10.4 dNTP mix (F-560)

The dNTP mix is a premixed ready to use solution consisting of the following compounds: dATP, dGTP, dCTP and dTTP dissolved in H₂O at 10 mM each.

10.5 50 mM MgCl₂ solution (F-510MG)

The 50 mM MgCl₂ solution can be used with the Mg²⁺-free DyNAzyme EXT Buffer or to supplement the Mg²⁺ concentration in the Optimized DyNAzyme EXT Buffer.

10.6 Lambda control template (F-304K)

The control template is bacteriophage lambda DNA (GenBank accession number J02459, 48 502 bp). The concentration is 0.5 ng/μl in TE-buffer.

10.7 500 bp control primer mix (F-557)

This component is a mix of primers for amplification of a 500 bp fragment of lambda DNA. Primer #1 is a 23-mer upper primer (5'-GAT GAG TTC GTG TCC GTA CAA CT -3') with a melting point of 64.4°C. The primer coordinates are 7131-7153 on the lambda DNA template. Primer #2 is a 23-mer lower primer (5'-GGT TAT CGA AAT CAG CCA CAG CG -3') with a melting point of 70.5°C. The primer coordinates are 7608-7630 on the lambda DNA template. Each primer concentration is 25 μM in H₂O.


10.8 20 kb control primer mix (F-558)

This component is a mix of primers for amplification of a 20 kb fragment of lambda DNA. Primer #1 is a 34-mer upper primer (5'-CTG ATG AGT TCG TGT CCG TAC AAC TGG CGT AAT C -3') with a melting point of 76.6°C. The primer coordinates are 7129-7162 on the lambda DNA template. Primer #2 is a 34-mer lower primer (5'-GTG CAC CAT GCA ACA TGA ATA ACA GTG GGT TAT C -3') with a melting point of 76.5°C. The primer coordinates are 27145-27178 on the lambda DNA template. Each primer concentration is 25 μM in H₂O.

10.9 Ready to use DNA standard (F-303SD)

This DNA standard is a mix of lambda DNA *Hind*III digest and bacteriophage φX174 DNA *Hae*III digest, each at 50 ng/μl (100 ng/μl total). It is supplied in 8 mM Tris-HCl (pH 8.0), 12 mM EDTA, 12 % glycerol and 0.012 % (w/v) bromophenol blue dye.

The DNA standard solution contains 19 fragments of the following sizes and mass amounts (per 10 μl):



Fragment	Base pairs	DNA amount ng/10 μl
1	23 130	238
2	9 416	97
3	6 557	68
4	4 361	45
5	2 322	24
6	2 027	21
7	1 353	126
8	1 078	100
9	872	81
10	603	56
11	564*	6
12	310	29
13a	281	26
13b	271	25
14	234	22
15	194	18
16a	125*	1
16b	118	11
17	72	7

Note: The cohesive areas of fragments 1 and 4 can be separated by heating to 65°C for 5 minutes. For daily use the marker can be stored at +4°C (at least 1 month). The marker is stable at -20°C for at least 1 year.

*Due to the low amount of DNA these bands are almost invisible.

10.10 Gel loading dye (F-350)

The gel loading dye contains: 50 % glycerol, 50 mM EDTA and 0.052 % bromophenol blue. For use, add 15 μl of gel loading dye to a 50 μl PCR reaction (or 6 μl to 20 μl PCR reaction).

10.11 Dimethyl sulfoxide DMSO, 100 % (F-515)

Note: The freezing point of DMSO is 18-19°C, so it does not melt on ice.

11. References

- Chester & Marshak (1993) *Analytical Biochemistry* 209, 284-290.
- Breslauer *et al.*, (1986) *PNAS* 83, 3746-3750.

Storage and shipping

DyNAzyme EXT PCR Kit is shipped on gel ice. Upon arrival, store the components at -20°C. DyNAzyme EXT PCR Kit is stable for one year from the packaging date when stored and handled properly.

Warranty

Finnzymes Oy warrants that its products will meet the specifications stated on the technical data section of the data sheets, and Finnzymes Oy agrees to replace the products free of charge if the products do not conform to the specifications. Notice for replacement must be given within 60 days of receipt. In consideration of the above commitments by Finnzymes Oy, the buyer agrees to and accepts the following conditions:

- That this warranty is in lieu of all other warranties, express or implied;
- That **ALL WARRANTIES OF MERCHANT ABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE ARE HEREBY EXCLUDED AND WAIVED**;
- That the buyer's sole remedy shall be to obtain replacement of the product free of charge from Finnzymes Oy; and
- That this remedy is in lieu of all other remedies or claims for damages, consequential or otherwise, which the buyer may have against Finnzymes Oy.

Exclusive terms of sale

Finnzymes Oy does not agree to and is not bound by any other terms or conditions, unless those terms and conditions have been expressly agreed to in writing by a duly authorised officer of Finnzymes Oy. Prices are subject to change without notice.

Recommended guidelines for safe use of the products

Finnzymes Oy recommends that the buyer and other persons using the products follow the N.I.H. guidelines published in the Federal Register, Volume 41, No. 131, July 7, 1976, and any amendments thereto. Finnzymes Oy disclaims any and all responsibility for any injury or damage which may be caused by the failure of the buyer or any other person to follow said guidelines.

Research use only

Since these products are intended for research purposes by qualified persons, the Environmental Protection Agency does not require us to supply Premanufacturing Notice.

Notice to user

The information presented here is accurate and reliable to the best of our knowledge and belief, but is not guaranteed to be so. Nothing herein is to be construed as recommending any practice or any product in violation of any patent or in violation of any law or regulation. It is the user's responsibility to determine for himself or herself the suitability of any material and/or procedure for a specific purpose and to adopt such safety precautions as may be necessary.

The quality system of Finnzymes Oy is certified according to standard SFS-EN ISO9001:2000.

DyNAzyme and UTW are trademarks of Finnzymes Oy or its affiliates.

DyNAzyme EXT DNA Polymerase is licensed under US Patent 5,416,149 owned by TaKaRa Shuzo Co.Ltd.

Version 2.5, May 2008